arXiv:quant-ph/0008106v1 25 Aug 2000

Off Resonant Pumping for Transition from Continuous to Discrete Spectrum and Quantum Revivals in Systems in Coherent States

G. S. Agarwal and J. Banerji
Physical Research Laboratory, Navrangpura, Ahmedabad 380 009, India

We show that in parametrically driven systems and, more generally, in systems in coherent states, off-resonant pumping can cause a transition from a continuum energy spectrum of the system to a discrete one, and result in quantum revivals of the initial state. The mechanism responsible for quantum revivals in the present case is different from that in the non-linear wavepacket dynamics of systems such as Rydberg atoms. We interpret the reported phenomena as an optical analog of Bloch oscillations realized in Fock space and propose a feasible scheme for inducing Bloch oscillations in trapped ions.

Keywords: Quantum revivals, Bloch oscillations, optical parametric interactions

I. INTRODUCTION

Parametrically driven systems have been studied extensively in quantum and nonlinear optics. Such systems have resulted in the production of twin beams of photons with high degrees of entanglement [1]. These beams of light have been used extensively in many applications such as in connection with Bell's inequalities and EPR paradox [2], quantum cryptography [3], teleportation [4], quantum imaging [5] and in general, in connection with the subject of quantum information processing [6]. The standard Hamiltonian describing such a parametric interaction is known to have continuous spectrum which then results in the exponential growth of (say) the energies in the signal and idler modes. The state of the system at any given time is, in fact, the so-called Perelomov coherent state [7].

In this communication, we report unusual features in the dynamics of systems in coherent states. We show how changing the detuning of the field pumping the system can cause a transition from a continuum energy spectrum of the system to a discrete one. This transition from continuum spectrum to discrete spectrum results in quantum revivals of the initial state.

This study has been motivated primarily by recent work on the optical realization of Bloch oscillations [8] and Wannier-Stark ladders [9]. These have been recently observed in ultracold atoms trapped in optical potentials [10], and in fields propagating in waveguide arrays with a linear bias so that the refractive index of a guide is proportional to its spatial location [11]. Even the effect of nonlinearities in the waveguide on Bloch oscillations has been studied. The underlying mathematical equation which leads to Bloch oscillations has the structure

$$i\dot{C}_n = n\Delta C_n + \beta(C_{n+1} + C_{n-1}) \tag{1}$$

In the absence of the term $n\Delta C_n$, an initial excitation in one of the C's gets delocalized , i.e, it spreads over all the other sites. However, if $\Delta \neq 0$, then the system returns to the original site. The linear dependence on the site index in the term C_n is very critical for the revival of the initial state. In the waveguide example, the linear bias produces the linear n dependence of the term $n\Delta C_n$. The optical realization of analogs of Bloch oscillations is based on situations where the dynamics could be described by equations like (1). We note that many coherently driven systems have dynamics characterized by equations like (1). Thus, remarkably enough, systems in coherent states can exhibit analog of Bloch oscillations. In the case of the radiation field, the index n would refer to the occupation number of a given Fock state. Thus the standard measurement of the photon number distribution can be used to study the analog of Bloch oscillations. Clearly this opens up the possibility of studying Bloch oscillations in a very wide class of driven systems. It is however important to have $\Delta \neq 0$, that is that the pumping should be off-resonant. Note also that the quantum revivals considered in this paper should be distinguished from the well studied quantum revivals in the nonlinear wavepacket dynamics of many systems such as Rydberg atoms [12] and coherent states propagating in a Kerr medium [13].

II. PARAMETRICALLY DRIVEN SYSTEMS

Consider a very commonly used system in optical physics. The Hamiltonian describing the parametric interaction of two modes a and b with frequencies ω_a and ω_b respectively, is given by

$$H = \hbar(Ga^\dagger b^\dagger + G^* ab). \tag{2}$$

Here G is the coupling constant which depends on the nonlinearity of the medium and the amplitude of the pump. It is assumed that the pump frequency $\omega_p = \omega_a + \omega_b$. The operators a and b satisfy Bosonic commutation relations. If the modes a and b are in vacuum state at time $t = 0$, then the state at time t is

$$|\psi(t)\rangle = \exp\{-it(Ga^\dagger b^\dagger + G^* ab)\}|0, 0\rangle \tag{3}$$

which is the so-called Perelomov coherent state or the squeezed vacuum state [7]. As is well known, the Heisenberg operator a, for example, evolves as

$$a(t) = a(0)\cosh(|G|t) - ib^\dagger(0)\frac{G}{|G|}\sinh(|G|t) \tag{4}$$

and thus the energy in the mode a continues to grow: $\langle a^\dagger(t)a(t)\rangle = \sinh^2(|G|t)$. The probability $p_0(t)$ of finding the system in the initial state is

$$\begin{aligned} p_0(t) &= |\langle 0,0|\exp\{-it(Ga^\dagger b^\dagger + G^* ab)\}|0,0\rangle|^2 \\ &= \cosh^{-2}(|G|t) \sim 4e^{-2|G|t}, \quad \text{as } t \to \infty. \end{aligned} \tag{5}$$

Thus the probability of returning to the initial state decreases monotonically and becomes exponentially small in the long-time limit (Fig. 1). Clearly, *there is no quantum revival* of the initial state.

The corresponding probability $p_n(t)$ $(n \neq 0)$ of finding the system in a state with n photons in each mode, is given by

$$p_n(t) = \cosh^{-2}(|G|t)\tanh^{2n}(|G|t); \quad n \neq 0. \tag{6}$$

Note that for $n \neq 0$, $p_n(t)$ increases with t initially and peaks at $|G|t = \tanh^{-1}\sqrt{n/(n+1)}$. The peak value of $p_n(t)$ has the expression

$$p_n(t)\rfloor_{\rm peak} = \frac{n^n}{(n+1)^{n+1}}, \tag{7}$$

which depends *only* on n and decreases monotonically with n.



FIG. 1. Evolution of a parametrically driven system when the pumping is resonant ($\Delta = 0$). The system dynamics is governed by the Hamiltonian given by Eq. (2). Here, $p_n(t)$ is the probability that the system, starting from vacuum, reaches a state at time t with n photons in each mode. Plots of $p_n(t)$ as a function of $|G|t$ are shown for $n = 0$, 1, 2 and 3. The plots are identified by their peak values which decrease as n increases. Note that the probability of returning to the initial state ($n = 0$) decreases monotonically with time.

The exponential growth of energies in the signal and idler modes and the subsequent absence of quantum revival for the initial state are connected with the fact that the Hamiltonian (2) admits a continuum of eigenvalues. The corresponding eigenstates are delta function normalizable and signify scattering states [14]:

$$\begin{aligned} H|\psi(\lambda)\rangle &= \lambda|\psi(\lambda)\rangle \\ \langle\psi(\lambda')|\psi(\lambda)\rangle &= \delta(\lambda - \lambda'); \quad \lambda,\ \lambda' \text{ real.} \\ |\psi(\lambda)\rangle &= \sum_n d_n|n,n\rangle, \end{aligned} \tag{8}$$

where the coefficients d_n are expressed as hypergeometric functions of the form ${}_2F_1(a,b;c,z)$.

Writing $|\psi(t)\rangle = \sum_n C_n(t)|n,n\rangle$ and substituting in the Schrodinger equation $i\hbar|\dot\psi\rangle = H|\psi\rangle$ with H as given in (2), we get

$$i\dot{C}_n = GnC_{n-1} + G^*(n+1)C_{n+1}. \tag{9}$$

Thus in terms of Fock states we have nearest-neighbor coupling with the coupling itself depending on the index n. The solutions (5)and (6) imply that with increasing time, an initial excitation in the $n = 0$ state spreads over all the states, or in the language of solid state physics, the excitation delocalizes.

We now consider an important physical situation where the Hamiltonian (2) is modified to

$$H = \frac{\hbar\Delta}{2}(a^\dagger a + b^\dagger b) + \hbar(Ga^\dagger b^\dagger + G^* ab). \tag{10}$$

This will be the case if the pump field is detuned, i.e, $\omega_p \neq \omega_a + \omega_b$. The amplitudes will satisfy Eq. (9) with an extra term $n\Delta C_n$ on the right. The state at time t can be obtained by using the general disentangling theorem [15] for the SU(1,1) operators $K_+ = a^\dagger b^\dagger$, $K_- = ab$ and $K_3 = (a^\dagger a + b^\dagger b + 1)/2$:

$$\begin{aligned} &\exp\{it(GK_+ + G^*K_- + \Delta K_3)\} \\ &= \exp(itG\Gamma_+K_+)\exp\{(\ln\Gamma_3)K_3\}\exp(itG^*\Gamma_-K_-) \end{aligned} \tag{11}$$

where

$$\begin{aligned} \Gamma_3 &= \left(\cosh\beta - i\frac{t\Delta}{2\beta}\sinh\beta\right)^{-2}, \\ \Gamma_\pm &= \frac{2\sinh\beta}{2\beta\cosh\beta - it\Delta\sinh\beta}, \\ \beta^2 &= -\beta_0^2 t^2, \\ \beta_0 &= \sqrt{\Delta^2/4 - |G|^2}. \end{aligned} \tag{12}$$

The probabilities are now given by $p_0(t) = |\Gamma_3|$ and $p_n(t) = |\Gamma_3||\Gamma_+|^{2n}$ where

$$\begin{aligned} |\Gamma_3| &= \left[1 + (|G|/\beta_0)^2\sin^2\beta_0 t\right]^{-1}, \\ |\Gamma_+|^2 &= \frac{\sin^2\beta_0 t}{\sin^2\beta_0 t + (\beta_0/|G|)^2} \end{aligned} \tag{13}$$

if the detuning Δ and the coupling constants are such that β_0 (defined by Eq. (12)) is real. Clearly the survival probability of the initial state is now an oscillatory function of time t and $p_0(t)$ is unity whenever $\beta_0 t$ is an integer multiple of π, i.e.

$$(\Delta^2/4 - |G|^2)t^2 = n^2\pi^2, \quad \text{where } n \text{ is an integer.} \tag{14}$$

That is, for these times, the system returns to the original state $|0,0\rangle$ up to an overall phase factor and there is a quantum revival (Fig. 2).



FIG. 2. As in Fig. 1 except that the pumping is off-resonant ($\Delta \neq 0$). The corresponding Hamiltonian is given by Eq. (10). Plots of $p_n(t)$ are shown for (a) $\Delta/2|G| = 1.01$ and (b) $\Delta/2|G| = 1.10$.

We thus find the important result that our parametric system exhibits analog of Bloch oscillation under the condition of non-zero detuning such that $\Delta > 2|G|$ and that the survival probability becomes unity for times defined by Eq. (14).

It is interesting to note that the peak values of $p_n(t)$ ($n \neq 0$) are still given by Eq. (7) although their locations, at $\beta_0 t = \sin^{-1}(\sqrt{n}\beta_0/|G|)$, will now depend on the detuning Δ.

Under the condition $\Delta > 2|G|$, the Hamiltonian (10) admits a discrete set of eigenvalues given by

$$\mathcal{E}_{qm} = \hbar\beta_0(q + 2m + 1) - \hbar\Delta/2; \quad m = 0,\, 1,\, 2,\, \ldots \tag{15}$$

where $q = a^\dagger a - b^\dagger b$ is an integer constant of motion. Thus the periodic behavior is connected with a transition in the nature of energy levels, i.e, the system has gone from a continuous spectrum to a discrete spectrum whose energy levels are equidistant. Note that the level separation is $2\hbar\beta_0$ which further validates our earlier observation that the revival period is given by $\pi/\sqrt{\Delta^2/4 - |G|^2}$ (see Eq. (14)). Comparison with the standard Bloch problem suggests that Δ plays the role of the electric field whereas the magnitude of G determines the coupling to the nearest neighbors.

III. OSCILLATOR DRIVEN BY A DETUNED COHERENT FIELD

All the above has analogs in the theory of coherent states for a *single* mode of the radiation field. In fact, coherent states will exhibit some new interesting physics if the detuning is nonzero. Consider the behavior of an oscillator driven by a detuned coherent field:

$$H = \hbar(\Delta a^\dagger a + \epsilon a^\dagger + \epsilon^* a). \tag{16}$$

If the system is in a vacuum state at $t = 0$, then the state at time t is a coherent state:

$$\begin{aligned} |\psi(t)\rangle &= \exp\left[\frac{i|\epsilon|^2}{\Delta^2}\left(\Delta t - \sin\Delta t\right)\right] \\ &\times \left|-i\epsilon\left(\frac{1-e^{-i\Delta t}}{i\Delta}\right)\right\rangle \end{aligned} \tag{17}$$

Consider first the case of zero detuning, $\Delta = 0$. In this case, the amplitude $-i\epsilon t$ of the coherent state continues to increase as t increases. The peak of the photon number distribution continues to move away from zero as t increases and the probability of being in the initial state decreases continuously and becomes exponentially small: $p_0(t) = \exp(-|\epsilon|^2 t^2)$.

For non-zero detuning, on the other hand, one obtains

$$\begin{aligned} p_n(t) &= |\langle n|\psi(t)\rangle|^2, \quad n \geq 0; \\ &= e^{-|\alpha|^2}\frac{|\alpha|^{2n}}{n!}, \quad |\alpha| = |\epsilon| t\left[\frac{|\sin(\Delta t/2)|}{\Delta t/2}\right]. \end{aligned} \tag{18}$$

Note that the peak of $p_n(t)$ occurs at $n = |\alpha|^2 = (2\epsilon/\Delta)^2\sin^2(\Delta t/2)$ which is an oscillatory function of time and vanishes whenever Δt is an integer multiple of 2π. At such times, $p_n(t) = \delta_{n,0}$, i.e, the system returns to the initial state $|0\rangle$ (Fig. 3) up to a phase factor. Note that the period does not depend on the strength of the driving field.



FIG. 3. Evolution of an oscillator driven by a coherent field of amplitude ϵ and detuning Δ. The corresponding Hamiltonian is given by Eq. (16). Plots of $p_n(t)$ as a function of $|\epsilon|t$ are shown for $n = 0$, 1 and 2 when (a) $\Delta/2|\epsilon| = 0$, (b) $\Delta/2|\epsilon| = 0.5$, (c) $\Delta/2|\epsilon| = 1.0$ and (d) $\Delta/2|\epsilon| = 2.0$. The plots are identified by their peak values $e^{-n}n^n/n!$, which decrease as n increases. All other notations are as in Fig. 1.

We also note that for $\Delta \neq 0$, the eigenvalues of the Hamiltonian (16) are equally spaced discrete states with energies

$$\mathcal{E}_n = n\hbar\Delta - \hbar|\epsilon|^2/\Delta. \tag{19}$$

The spacing is independent of ϵ which leads to a periodic behavior independent of ϵ. On the other hand, if $\Delta = 0$, then $H = \hbar(\epsilon a^\dagger + \epsilon^* a)$ has a continuum of eigenvalues with the eigenfunctions satisfying the usual delta function condition $\langle\psi(x)|\psi(x')\rangle = \delta(x - x')$. Thus here again we see the transition from continuum eigenvalues to the discrete set [16] and this transition is responsible for the revival of the initial state.

IV. QUANTUM REVIVALS IN ION TRAPS

We now show that this simple behavior can be easily probed, as for example, for a trapped ion like the one used by Wineland and co-workers [17]. Consider an ion in a deep trap cooled to its ground state. Let it interact with two Raman fields E_1 and E_2 with frequencies ω_1 and ω_2 such that $\omega_1 - \omega_2 - \nu \neq 0$ (Fig. 4).



FIG. 4. Raman-like off-resonant transitions among the vibrational states of an ion cooled to its electronic ground state in a trap. The ion is acted upon by two fields with amplitudes E_1 and E_2 and frequencies ω_1 and ω_2 respectively. The separation between the relevant vibrational levels is denoted by ν so that the detuning $\Delta = \omega_1 - \omega_2 - \nu$. The ion would return to its ground state whenever $\Delta t = 2n\pi$, where n is an integer.

In such a case the effective Hamiltonian describing the ionic motion will be of the form $E_1 E_2^* a \exp\{i(\omega_1 - \omega_2 - \nu)t\} + h.c.$ This Hamiltonian is equivalent to (16). Here a is the annihilation operator for the vibrational motion of the ion. A measurement of the vibrational distribution will exhibit the analog of Bloch oscillation provided $\omega_1 - \omega_2 - \nu \neq 0$ and the ion would return to its ground state whenever $(\omega_1 - \omega_2 - \nu)t = 2\pi n$, where n is an integer.

V. CONCLUSION

In conclusion, we have introduced off-resonant pumping of coherent systems as a new mechanism for the optical analog of Bloch oscillations. In our scheme, such oscillations occur in Fock space. We showed that for parametrically driven systems and systems in coherent states, the detuning of the pump changes the energy spectrum of the system from continuum to discrete , and revives the initial state. Finally, we proposed a scheme for inducing Bloch oscillations in trapped ions.

---

[1] Greenberger D M Horne M A and Zeilinger A 1993 Physics Today **xx** 22

[2] Ghosh R and Mandel L 1987 Phys. Rev. Lett. **59** 1903

[3] Ekert A K 1991 Phys. Rev. Lett. **67** 661

[4] Bouwmeester D, Pan J-W, Mattle K, Eibl M, Weinfurter H and Zeilinger A 1997 Nature **390** 575; Boschi D, Branca S, Martini F De, Hardy L and Popescu S 1998 Phys. Rev. Lett. **80** 1121

[5] Kolobov M I 1999 Rev. Mod. Phys. **71** 1539

[6] Bennet C H and Shor P W 1998 IEEE Trans. Inf. Theory **44** 2724

[7] Perelomov A M 1986 *Generalized Coherent States and Their Applications* (Berlin: Springer); Caves C M and Shumaker B L 1985 Phys. Rev. A **31** 3068, 3093

[8] Bloch F 1928 Z. Phys. **52** 555; Zener C 1932 Proc R Soc London A **154** 523

[9] See for example Wannier G H 1959 *Elements of Solid State Theory* (London: Cambridge University Press) pp 190-193; For a recent optical realization of Wannier-Stark ladder see de Sterke C M, Bright J N, Krug P A and Hammon T E 1998 Phys. Rev. E **57** 2365

[10] Dahan M B, Peik E, Reichel J, Castin Y and Salomon C 1996 Phys. Rev. Lett. **76** 4508; Wilkinson S R, Bharucha C F, Madison K W, Niu Q and Raizen M G 1996 *ibid* **76** 4512

[11] Morandotti R, Peschel U, Aitchison J S, Eisenberg H S and Silberberg Y 1999 Phys. Rev. Lett. **83** 4756; Lenz G, Talanina I and de Sterke C M 1999 *ibid* **83** 963; Pertsch T, Dannberg P, Elflein W, Bräuer A and Lederer F 1999 *ibid* **83** 4752 ; Peschel U, Pertsch T and Lederer F 1998 Opt. Lett. **23** 1701

[12] See, for example, Averbukh I Sh and Perelman N F 1989 Phys. Lett. A **139** 449; Gaeta Z D and Stroud Jr C R 1990 Phys. Rev. A **42** 6308

[13] See for example Yurke B and Stoler D 1986 Phys. Rev. Lett. **57** 13; Tara K, Agarwal G S and Chaturvedi S 1993 Phys. Rev. A **47** 5024

[14] Agarwal G S and Puri R R 1996 Int. J. Mod. Phys. **B10** 1563

[15] Wódkiewicz K and Eberly J H 1985 J. Opt. Soc. Am. B **2** 458; Barnett S M and Radmore P M 1997 *Methods in Theoretical Quantum Optics* (Oxford: Clarendon Press)

[16] We do not consider systems belonging to the group SU(2) as their spectrum is discrete anyway and the system would always exhibit periodic behavior due to the compactness of the SU(2) group

[17] Meekhof D M, Monroe C, King B E, Itano W M and Wineland D J 1996 Phys. Rev. Lett. **76** 1796; Leibfried D, Meekhof D M, King B E, Monroe C, Itano W M and Wineland D J 1996 *ibid* **77** 428